

Magnetic Correlates in Electromagnetic Consciousness

A R Liboff, Department of Physics, Oakland University, Rochester, MI

Abstract. We examine the hypothesis that consciousness is a manifestation of the electromagnetic field, finding supportive factors not previously considered. It is not likely that traditional electrophysiological signaling modes can be readily transmitted throughout the brain to properly enable this field because of electric field screening arising from the ubiquitous distribution of high dielectric lipid membranes, a problem that vanishes for low frequency magnetic fields. Many reports over the last few decades have provided evidence that living tissue is robustly sensitive to ultrasmall (1-100 nT) ELF magnetic fields overlapping the γ -frequency range often associated with awareness. An example taken from animal behavior (coherent bird flocking) lends support to the possibility of a disembodied electromagnetic consciousness. In contrast to quantum consciousness hypotheses, the present approach is open to experimental trial.

KEY WORDS: Electromagnetic Consciousness, Quantum Consciousness, ELF Magnetic Field, Bird Flocking, Mind Sharing

Introduction. The great advantage in any field approach to the problem of consciousness is its unifying character, allowing one to regard consciousness in a global manner (Pockett, 2013). More specifically, recent arguments have suggested (Pockett, 2000, 2012, 2013; McFadden, 2001, 2002, 2007, 2013) that the electromagnetic (EM) field may play a unifying role in brain function, facilitating the binding of multimodal sensory data viewed as integral to conscious experience. Pockett (2000), focusing on the extensive appearance of seemingly independent oscillatory neural processes, has proposed the provocative hypothesis that *“Consciousness is identical with certain spatiotemporal patterns in the electromagnetic field”*. There is no question that incorporating electromagnetic field-mediated communication into models of brain function has the potential to reframe discussions surrounding consciousness. However to the best of our knowledge all previous discussions concerning this possibility have only focused on traditional electrophysiological parameters, with no recognition of the role likely played by the magnetic field.

Electromagnetic fields generated within localized brain structures can exert effects more globally within the brain and may therefore transcend purely connectionist models of brain function. Related phenomena involving interactions generated from multiple sources, particularly those involving the frequency, phase and coherence of oscillatory fields, suggest a more unitary view of brain function often regarded as an important requirement for consciousness. Such field mediated communication within the brain suggests that the laws of physics pertaining to electricity and magnetism and signal processing may be more applicable in this context than had previously been supposed.

In considering the nature of such electromagnetic field-mediated communication within the brain, one must recognize that there are severe physical constraints involved. The voltages and currents generated during neural excitations are extraordinarily weak, so much so that important questions arise as to limits on detectability. In partial response to this question of signal threshold McFadden (2001, 2002) pointed out that the case for an electromagnetic contribution to brain function and consciousness is strengthened considerably when one considers increases in field intensity resulting from synchronicity, i.e., many neurons acting in concert. However no evidence is provided to indicate how such synchronicity is enabled.

In the following, we explore general questions related to electromagnetic field-mediated communication within the brain and its relationship to consciousness and investigate field levels possible given the physical characteristics of the nervous system. We broaden the focus that limits consideration to traditional electrochemical strategies, widening the possibility to include ultraweak magnetic fields derived from axonal and post-synaptic charge flows. A more comprehensive literature-based provenance of experimental studies is provided detailing electromagnetic effects on the brain. Finally, we point to one interesting potential consequence of using a field concept to describe consciousness, applying a gedanken model to cooperative animal behavior to illustrate a primitive electromagnetic neural scenario.

Characterizing the Electromagnetic Field. The electromagnetic (EM) field, similar to the other three fields in nature (gravitational, strong, and weak) is defined in terms of the force it exerts. Thus, the intensity of the EM field at any point in space is measured by the force it exerts on electric

charges and magnetic dipoles. Because of its vector nature, its direction must be specified at each point. A field can be static, i.e., unchanging in time, or time-varying. It can be uniform such that within a given region it is the same in intensity and direction, or it can vary from point to point. These differences are determined by the sources of the field, the electric charges or the magnetic dipoles or both. When we attempt to relate the electromagnetic field to the brain and specifically to its possible role in consciousness, we can anticipate that any such field must be derived from both electric and magnetic sources, that it will be quite non-uniform, and that it must necessarily vary in time.

In its most literal sense, there is only one EM field, one that fills all of space. This should not be taken to mean that electromagnetic consciousness, if it truly exists, implies some sort of connecting link between *all* conscious creatures. The presumption of information transfer underlies EM consciousness, and although the EM field extends to great distances, its ability to convey information falls off rapidly with distance due to decreasing signal-to-noise ratios.

The precise mathematical understanding of the electromagnetic field dates back to its formulation by James Clerk Maxwell in the latter half of the 19th century. Perhaps the most important thing about Maxwell's Equations is the way they reveal the interconnectedness of the electric and magnetic vector components that comprise the EM field. These components are often labeled E and B, respectively, for electric field and magnetic field. The electric field strength in the brain is mostly manifested by the voltage differences across cell membranes. The magnetic field is determined by currents, derived from the passage of charge along axons and associated neural tissues. This distinction between electric and magnetic sources is mirrored when discussing action potentials as opposed to current dipoles. A similar dichotomy occurs when comparing EEG to MEG, as something solely dependent on E and something solely dependent on B. In making this distinction the reader is cautioned against thinking that these two measures of the electromagnetic field are merely two sides of the same coin. The electric properties of the brain are quite different from the magnetic.

To properly describe the electromagnetic field in any substance it is necessary to know the electrical and magnetic properties of that substance. The two key parameters used to characterize different substances, including brain tissues, are the electric permittivity ϵ and magnetic permeability μ . With very few exceptions the permeability of all living tissues, including that

of the brain, is hardly different from free space. This means that neural tissues are transparent to magnetic fields. On the other hand the permittivity ϵ varies considerably from point to point in the brain, mostly reflecting the large differences in conductivity among lipid membranes, axonal fluids, and neurotransmitters.

This fact makes it difficult to conceive of a mechanism by which information, in the form of electric field changes, could be successfully transported from within an individual cell body to remote areas in the brain. The difficulty lies with the cell membrane, whose E-field is so large that it considerably reduces the external transfer of electric-field changes generated within the axon or dendrite, an effect that can be thought of as membrane shielding. ~~Indeed it is generally recognized that one of the important functions served by lipid membranes is to protect the cell interior from unwanted external electric field effects. It is equally true that this same membrane shield prevents electric signals within the cell from being easily transmitted. Instead throughputs in the form of ion channels and pumps are used to provide transmembrane information flow.~~

This problem is not shared with magnetic fields. The magnetic field generated by a current-carrying axon is in principle capable of being detected throughout the cranium, the only limitation being the size of the current. This distinction does not mean that any electromagnetic field representing consciousness is necessarily independent of the local electric field. The E field within neural tissue is probably not ~~electrically involved in responsible for~~ information transfer over large regions of the brain ~~as implied in Pockett's hypothesis~~, but it is likely ~~serves involved in EEG and similar volume conduction effects. an enabling role by which the transport of electric charge is made possible.~~

Even though a complete description of the electromagnetic field is dependent on knowing both E and B, there are situations where the EM field is adequately described simply as either an E-field or as a B-field. For example the electromagnetic field in a region where there are no moving charges is purely an electric field. Conversely, B-fields occur where one is only concerned with the presence of magnetic dipoles or effects due to moving charges, with the local electric field either unimportant or very small. This is particularly important within the range of the extremely low frequencies (ELF) that might be associated with consciousness, i.e., below a few hundred Hz. Indeed, there is good reason to believe that weak ELF

magnetic field effects may be an essential component of the connection between the electromagnetic field and consciousness.

Mind as an Expression of Quantum Theory. Various speculations have been advanced (Hameroff, 1994, 2012; Penrose, 1989, 1994; Hameroff and Penrose, 2014; Baars and Edelman, 2012) that consciousness has its basis in emergent physics, specifically the remarkable observations (Aspect et al, 1982) which upset our notions about particle locality, summarized in the phrase quantum entanglement. Well-accepted features of reality in physics such as nonlocality and an energy-filled vacuum are also potentially attractive concepts for use in consciousness. In some respects this approach to consciousness is an extension of the older Copenhagen school of quantum mechanics, originating with Bohr and Heisenberg, that regarded mind as an ingredient of reality, realized when an individual makes a choice that collapses one wave function to the exclusion of all others. In another respect the involvement of quantum theory can be seen as conflating computation with consciousness, the argument boiling down to saying that thinking is the same as large-scale parallel computing. In any event it has proven difficult to conceive of an experiment to test this idea. Perhaps the most telling criticism of the quantum approach to consciousness came from Chalmers, who proposed “... a Law of Minimization of Mystery: consciousness is mysterious and quantum mechanics is mysterious, so maybe the two mysteries have a common source” (Chalmers, 1995).

Worth noting in the Hameroff/Penrose quantum approach is the reliance on qubits, abstract elements used in quantum computing that are assumed to be associated with microtubules. But in so doing the idea of a field approach is compromised amounting to an attempt to generalize from a reductionist base, a process hardly different from the usual neurochemical approaches to consciousness. In sharp contrast the electromagnetic field explanation begins with the broader statement rather than the detailed.

Certainly, quantum physics plays an important role at the atomic level in living things. The energy levels in an uncoupled hydrogen atom found in a mitochondrion are no different, quantum-wise, from those in hydrogen atoms found anywhere in inanimate matter. What matters in living things, however, is the configuration of all the mitochondria in a single system assembled according to the electromagnetic field. In short we are of the opinion that quantum mechanics, as good as it is with microscopic details, is an

~~inappropriate means of looking at the organizational side of things, whether considering mind or life itself.~~

~~Quantum mechanics is not the only way physics can help in explaining consciousness. We draw attention to the idea, expressed in greater detail below, that the electromagnetic field as expressed in the brain may represent only one specific portion of the overall electromagnetic field associated with the complete organism. This brings into question the need for a separate, independent application of quantum theory to consciousness. One might think that theories about consciousness from the world of physics should specifically reflect overall biological function or relate to the living state *per se*. It might be supposed that particle entanglement, given its profound nature, is necessarily also involved in the question of life itself. Microtubules, figuring so prominently in Hameroff/Penrose thinking, are not confined to the brain, but are found in all tissues.~~

The Electromagnetic Origin of Mind. Therefore, Quite apart from limiting consideration solely to the brain, the nature of biological expression itself may be essentially electromagnetic, something previously theorized by the author (Liboff, 1994, 2004, 2005). If this is the case then it is reasonable to think that the brain, as one part of the entire organism, should also reflect this property.

In this scenario life is regarded as a consequence of the electromagnetic field of force, one of the four force fields that physics uses to describe reality. The living state, in this view, falls naturally into the increasingly complex sequence of electromagnetic configurations that nature has constructed beginning first with the simple delineation of electric charge in electrons, through successively higher-level systems of atoms, then molecules, then polymers, and finally into the present electromagnetic configuration that we describe as the living state. In this context, just as the appearance of atoms and molecules in nature is hardly mysterious, this is no less true for life.

This categorization of living things as an expression of the electromagnetic field still manages to reflect the usual cast of biological/biochemical observables that are found genetically, somatically and instantaneously. In this view all the detailed biochemical parameters that comprise the entire organism-the component enzymes, lipids, DNA/RNA states, sugars, etc, - are regarded as contributing to an equivalent electromagnetic polarization vector that varies in time as the organism changes. The biochemical details

are maintained exactly as described in present-day textbooks, but their effects are replaced by the global field that results from the overall instantaneous configuration of charges and currents. Using this approach one level of reductionist detail is removed and an overarching generalization is revealed. The EM field viewed in detail is no less complex than the components it replaces, but as a whole it represents a unifying synthesis, dependent only on electromagnetic parameters such as E , B , ϵ , and μ .

Given this approach to synthesizing the biological components of the entire living thing, one cannot reasonably exclude the brain, as part of the whole, from also being described in similar fashion. One can therefore perhaps regard consciousness as merely one more example of electromagnetic biology, something uniquely specific to the brain. In this view associating the electromagnetic field with brain function does not mean that this field is anything more than a convenient description, merely corresponding to what is observed elsewhere in the overall biological system. Nonetheless, the interesting question remains: If the field description of living things is an accurate measure of the living state does this imply that it also carries the role of biological expression, that the living state unfolds in time as dictated by the EM field? This has the effect of evolution “capturing” the EM field, in the same way as other physical realities are utilized, absorbed into the genome for purposes of survival. If one adopts this point of view then it can also be argued that consciousness is an expression of the electromagnetic field. Just as the body’s EM field helps maintain liver function and homeostasis, it also serves to realize brain function.

In any event, regardless of how electromagnetic consciousness came about, whether as part of something larger, or as a unique addition, the outcome had enormous evolutionary significance, enabling the nervous system to not only react to stimuli, but also to control them. We imagine evolution as recognizing the electromagnetic field as a powerful adaptation for the nervous system, providing higher level animals and particularly humans with a unique survival strategy.

Weak Extremely Low Frequency (ELF) Magnetic Interactions. The neurophysiological community has failed to take note of the abundant peer-reviewed literature reporting that weak ELF magnetic fields interact with living things. Many of these reports indicate that this type of interaction is observed at remarkably small intensities, a fact that has relevance to the

general question of electromagnetic consciousness. One cannot simply ignore this literature base, as when McFadden (2002) wrote “*The issue of the sensitivity to weaker [fields] is entangled with the power line controversy, which despite many studies, remains contradictory and unresolved.*” It may be true that the power line controversy, primarily epidemiological in nature, remains unresolved, but a host of well-replicated independent laboratory studies (Liboff et al, 1984; Goodman and Henderson, 1988; Walleczek and Liburdy, 1990; Fitzsimmons et al, 1995; Liboff, 2006) reveal that the interaction capabilities of weak magnetic fields with living tissues are reasonably certain.

By far the most significant of these was the work by Zhadin et al (1998) reporting that AC magnetic fields as weak as 40 nT can significantly affect the conductivity of glutamic acid in aqueous suspension, an observation not only consistent with earlier animal studies (Martin and Lindauer, 1977; Walker and Bitterman, 1989; Beason, 2005) but later replicated elsewhere independently (Pazur, 2004; Comisso et al, 2006; Alberto et al, 2008). This result, involving vanishingly small magnetic intensity opens the door to serious consideration that magnetic fields may play a heretofore unsuspected role in brain function. Even earlier, considerable evidence had accumulated showing that neural systems are sensitive to weak magnetic fields, particularly in studies on rat behavior (Thomas et al, 1986; Lovely et al, 1993; Lyskov et al, 1996; Zhadin et al, 1999). The primary conclusion of these studies was that activation of the N-methyl D aspartate (NMDA) glutamate receptor, deeply involved in synaptic plasticity, memory, and learning, is affected by weak ELF magnetic fields (Manikonda et al, 2007; Frilot et al, 2014). In this connection note that Flohr (2000) argues that the plasticity of NMDA receptors is a necessary condition for consciousness.

Further, work on anesthetized rat hippocampus has shown that electrical activity among pyramidal neurons is affected by weak ELF magnetic fields (Jenrow et al, 1998). These magnetically-induced changes persisted for tens of minutes after removal of the field, lending strength to the likelihood that such fields affect synaptic plasticity. The investigators in this work also hypothesized the existence of endogenous magnetic fields “mediating” neuronal synchrony. To add to this picture, other studies have reported the EEG output in rats and in human is significantly affected by ELF weak magnetic fields, applied either as pure sinusoidal waves or as modulations of higher radiofrequency signals (Takashima, 1979; Vorobyov et al, 1997; 1998; 2010; Marino et al, 2004). Finally, in studies that may have no bearing on consciousness *per se*, but nevertheless lend strength to the remarkable

sensitivity of neural tissue to magnetic fields, various investigators have found that proliferation in neuroblastoma cell culture is also affected by weak ELF magnetic fields (Smith et al, 1992; Horton et al, 1993; Blackman et al, 1994).

This large body of literature impacts the question of electromagnetic consciousness. If, **as so thoroughly demonstrated in these reports**, externally applied weak magnetic fields are interactive with neural tissue, then the endogenous equivalent of such fields must also be considered as possible factors involved in consciousness.

Synchronicity. The question of synchronicity arises when trying to demystify binding, that aspect of consciousness whereby long-range seemingly simultaneous recruitment of multiple sensory inputs occurs, enabling one to experience the unified sense of being (Jefferys et al, 1996; Tomoni and Edelman, 1998). No satisfactory explanation has yet been agreed upon. Synchronicity can be approached in **two various** ways, **not the least of which is Adey's vision of cells "whispering together"**. **First,** There is the potential for enhanced information transfer when action potentials fire simultaneously, either among parallel arrays of axonal fibers or by means of some remote coupling processes. Presumably, this type of synchronicity is subject to traditional electrophysiological constraints, including the relatively long times associated with biochemical signal creation at cell membranes. A single axonal voltage change can be used to advantage by the many post-synaptic connections, leading to large-scale dissemination of information throughout the brain. However it is important to note that the coupling mechanism, whereby multitudes of like processes occur synchronously, remains obscure.

The second Another type of postulated synchronicity is long-range resonance recognition, where remote areas of the brain appear to be sensitive to specific frequencies referred to as gamma waves, often occurring in the 25-100 Hz range. Considerable weight is attached to the observation that gamma waves disappear under anesthesia, tending to reinforce the notion that they are a measure of awareness. We find good reason to think that endogenously generated magnetic fields are directly related to the phenomenon of synchronicity.

This range of gamma wave frequencies is remarkably close to the frequency range that is found in the various ELF studies mentioned above. Of particular

interest are those interactions occurring at ion cyclotron resonance (ICR) frequencies (Liboff, 2006). Not only do gamma wave frequencies correspond closely to these ELF interactive frequencies, but the overall timing is greatly reduced when comparing magnetic field changes to electrophysiological processes. Most critically, the well-described resonance capabilities of living things to specific magnetic frequencies (Liboff, 1985; Fitzsimmons et al, 1995; Zhadin et al, 1998, 1999; Vorobyev et al, 1998; Baureus-Koch et al, 2003; Gaetani et al, 2009; Foletti et al, 2013) provides a reasonable basis on which to begin to explain large-scale synchronicity.

Also, because of membrane “shielding” we specifically rule out any global effects due to long-range cooperative electric field changes. It has already been pointed out that the neural environment includes a wide variety of electrical conductivities, making it difficult to retain or distribute coherent systematic changes in electric field. Indeed **as mentioned above** it has often been observed that one important function of the cell membrane is to shield the interior against unwanted external electrical perturbations. This natural barrier to *electric* communication does not similarly apply to remote *magnetic* signals generated by axonal currents. Thus, although the view of an electromagnetic spatiotemporal structure to express consciousness has great merit, it is unlikely that the electric field vector E alone is capable of providing a suitable means of information transfer through synchronicity. On the other hand, any magnetic field component, generated locally, has wide access to the entire brain, particularly at low frequencies. In short that part of the brain’s EM field related to binding is due to oscillatory magnetic fields that are derived from local electric currents. In this way, complex information transfer may be enabled throughout the brain.

It is interesting that neuroscientists do not ordinarily view the information transfer that occurs during neural excitation in terms of electric current. In part this is because ionic displacement is involved, as opposed to electron flow. Further, the most readily available experimental variable is the voltage change at the scalp, as measured in EEG. Even the classic squid axon preparation routinely categorizes propagation signals in terms of voltage. One refers to evoked potential, rarely to evoked current. But the fact remains. Potential differences may be easier to measure, but the critical event necessarily involves charge transfer. The most elementary neurophysiological event is always the result of a change in net charge δq , occurring over the time δt , giving rise to a current $i = \delta q / \delta t$, and therefore by virtue of the Biot-Savart Law, an associated magnetic field. The complex of such individual mini-fields in a single brain can be summed over their

intensities and relative directions to result in an *uberfeld* that is both unique to the individual and a physical realization of his/her consciousness.

It is useful to estimate the level of current required to produce a detectable magnetic field. Reliable reports (Zhadin et al, 1998) discovered that fields as small as 10-100 nT and, in one case, between 1 and 10 nT (D' Emilio et al, 2014) can be detected in biological and biochemical systems. Accordingly, we will assume that in order for a magnetic field to be found useful in the nervous system, its intensity must be on the order of 1-100 nT. However, in order to achieve a basis for electromagnetic consciousness, not only is it required that neural elements are capable of detecting the consequent magnetic information, but such elements must also be capable of generating this information. It is therefore critical to determine the maximum possible magnetic intensity generated by current elements, presumably of axonal and dendritic origin. It is tempting to believe that fields of 1-100 nT are endogenously generated in the brain, something that would provide a strong likelihood for synchrony and binding. In this connection we note that this range of magnetic intensity has yet to be explored. Magnetoencephalography (MEG) has specifically focused on much smaller field intensities, ranging from 10s of fT (10^{-15} T) to 1000s, weaker by a fraction of about ten thousand than our projected values. However MEG devices are specifically designed to ~~provide electronic limitations on~~ **prevent the measurement of** unwanted larger biomagnetic signals, certainly those in the 1-100 nT region.

At much larger magnetic field intensities, the lack of any perceived neurological effects resulting **from** Magnetic Resonance Imaging (MRI) has also led some to discount the possibility of electromagnetic consciousness. However, one distinguishing feature of the observed responses to ELF magnetic fields has been the fact that these responses are not found at higher intensities, but limited to certain weak intensity **levels windows** (Bawin et al, 1976; Blackman et al, 1989; Lednev, 1991), in all cases many orders of magnitude less than MRI intensities. Indeed recent studies indicate that these intensities are so small as to warrant the use of hypomagnetic research facilities (Pazur, 2004; Comisso et al, 2006; Alberto et al, 2008, D' Emilio et al, 2014), **without which, as first observed by Zhadin (1998) the effects of these vanishingly small fields are not detectable.**

Can Consciousness be Shared? We have already mentioned that the transfer of useful electromagnetic information falls off rapidly with distance from the source of the field. However it must be noted that this falloff can be finessed to some extent by signal processing. Very narrow frequency bandwidths can in some cases be detected in the presence of noise, providing that source and receiver are properly matched in frequency response. If indeed EM consciousness is a reality, involving very weak magnetic fields, this could have interesting consequences. For example some fraction of one's mental field might be available to others nearby. We can offer no evidence in support of this possibility. Nonetheless, considering the profound nature of what is involved, it is worth examining the potential effects on human interactions.

In approaching this question, we focus on a long-standing problem in animal behavior, the remarkable cooperative interaction found during bird flocking (~~Fourie and Greenwold, 2002;~~ Cavagna et al, 2010; Hemelrijk and Hildenbrandt, 2011), a phenomenon that can involve tens of thousands of birds at a time, and often extends ~~ing~~ over the greater fraction of a kilometer. ~~One puzzling mystery is that the correlation between birds does not decay with distance (Cavagna et al, 2010), making it difficult to believe that information is can be conveyed as cues from bird to bird in to enable what appears to be a unitary response. The main problem with trying to explain this strange effect is the speed with which We can immediately reject the notion that bird-to-bird communication is conveyed by cues. There simply is not sufficient time to transmit and receive visible signals at distances of perhaps 0.5 km so quickly as to appear instantaneous. Instead~~

We raise the possibility that neuroelectromagnetic signals originating in ~~the a~~ bird's brain ~~traveling at the speed of light~~ may serve as a means of nearly instantaneous, speed of light, information transfer among all the birds. ~~That is, Instead of bird-to-bird sensing of visual cues—how likely is it that~~ an electromagnetic field is, in effect, shared by the entire flock.

We can attempt to flesh out such ~~this type of~~ behavior in terms of ~~a~~ this commonly sensed electromagnetic field. Consider a large flock of birds, each of which is equally capable of both generating a local field as well as responding to such a field. Assume that all the individuals comprising this flock can equally transmit and receive the same neural information, conveniently labeled as $f_i(B)$, where the index i specifies the individual bird

and where the functional dependence is on the local magnetic field B . We can therefore state that $f_i(B) = f_{i+1}(B)$ for each and every member of the group. A little thought will reveal that this function can also be replaced by a single representation for the entire group. If we arbitrarily associate $f_i(B)$ with the movement of the i th bird this connection to movement must also apply to all the birds in this flock. Each bird contributes to the whole, but for a large flock adding or subtracting one or even a few birds does not affect the bulk activity of the larger entity comprising the flock. Also, more than merely the magnetic intensity alone would be needed to provide bird-to-bird signaling. Without knowing precisely how the function $f(B)$ is constructed, it seems clear that it must reflect a *change* in B , rather than B itself.

In one sense this provides a *de facto* explanation, however speculative, of all cooperative motional phenomena such as bird flocking and fish shoaling. But a separate conclusion can also be drawn. Simply stated this cooperative neural behavior arising from the summation of the many individuals in the flock is indistinguishable from that of a single imagined system exhibiting an equivalent unitary behavior. **Indeed Couzin (2007) deliberately used the phrase “collective mind” to describe this behavior. In this** We add to this view, **suggesting that** the contribution of the individual bird vanishes, and is replaced by a unique “nervous system”, common to all. In effect we idealize the neural characteristics of the entire group in terms of a single entity that represents the whole. This entity, although determined from the neural activity of individual birds, is nonetheless independent of individual birds.

This entity amounts to an electromagnetic distillate of neural function, illustrating in a very basic way how an electromagnetic field can be thought of as something that describes brain activity but is still independent of its individual elements. We make the leap to electromagnetic consciousness, arguing that it too can be regarded as a stand-alone physical entity.

Discussion. Various speculations have been advanced (Hameroff, 1994, 2012; Penrose, 1989, 1994; Hameroff and Penrose, 2014; Baars and Edelman, 2012) that consciousness has its basis in emergent physics, specifically the remarkable observations (Aspect et al, 1982) which upset our notions about particle locality, summarized in the phrase quantum entanglement. Well-accepted features of reality in physics such as nonlocality and an energy-filled vacuum are also potentially attractive concepts for use in consciousness. In some respects this approach to

consciousness is an extension of the older Copenhagen school of quantum mechanics, originating with Bohr and Heisenberg, that regarded mind as an ingredient of reality, realized when an individual makes a choice that collapses one wave function to the exclusion of all others. In another respect the involvement of quantum theory can be seen as conflating computation with consciousness, the argument boiling down to saying that thinking is the same as large-scale parallel computing. In any event it has proven difficult to conceive of an experiment to test this idea. Perhaps the most telling criticism of the quantum approach to consciousness came from Chalmers, who proposed “... *a Law of Minimization of Mystery: consciousness is mysterious and quantum mechanics is mysterious, so maybe the two mysteries have a common source*” (Chalmers, 1995).

We have shown that physics can play a central role in the problem of consciousness, quite independently of the approach associated with quantum modeling. In particular there is good reason to support a modified version of the Pockett hypothesis that consciousness is a consequence of the electromagnetic field. Specifically, we argue that endogenous low-frequency magnetic signals derived from axonal and neurotransmitter charge flow must also be considered in addition to the more readily recognized electric parameters. Our argument is strengthened by the many reports over the past few years showing that neural expression is surprisingly responsive to weak magnetic fields, particularly in the 25-100 Hz frequency range implicated by many with the question of binding, lending one to believe that such fields are similarly generated in the brain.

Without the presence of such fields, it is not clear that a reasonable electromagnetic hypothesis is possible. There simply is no obvious means of large-scale, electrophysiological information transfer from axonal sources to remote areas in the brain. This is because the ubiquity of lipid membrane structures in the brain ensures that axonal voltage changes are not directly shared externally. Lacking a magnetic means of communication the global delivery required for unitary consciousness instead reverts to the present biochemical paradigm, which is, by definition, completely different from a field model.

We envision electromagnetic consciousness as necessarily requiring the capability of rapid *intra-brain* information transfer. In order to properly utilize the electromagnetic field this requirement carries important physical constraints. Thus, electromagnetic consciousness is dependent both on the level of sensitivity to very weak fields within neural tissues as well as the

corresponding capability to also generate detectable currents. Our present knowledge indicates an approximate sensitivity of 1-100 nT within biological tissues, suggesting that there is good reason to search for endogenously generated magnetic signals in this range. Unlike quantum explanations of consciousness, ~~the—~~ **an** electromagnetic hypothesis involving these field intensities can ~~readily~~ be experimentally explored.

The possibility that such magnetic signals at these intensities are generated by the brain implies something not possible with the present neuroelectric paradigm, namely that signals originating within the brain may be detected externally by others, suitably attenuated. Thus, if consciousness is an expression of the electromagnetic field, then one possible consequence may be near-field *interbrain* communication. This should not be construed as akin to extrasensory perception, but rather as a previously undescribed type of neurological effect perhaps best imagined as mind sharing. It is unlikely that the individually detailed complex gestalt that constitutes one's consciousness field can be successfully transmitted in coherent form to external systems. Rather we are suggesting that limited fractions of an individual's electromagnetic field are sufficiently strong to bridge the neurological activity in nearby sentient systems, possibly appearing as unrecognized magnetic cues. This distinction between subtle cues and full-blown thoughts can be taken to mean that even if consciousness per se cannot be electromagnetically shared, there may still be weakly interactive influences. We may be immersed in such magnetic cues without being aware of it.

An interesting speculation follows from the two sharply contrasting neural approaches, one the presently accepted biochemical/electrophysiological paradigm, the other involving the electromagnetic field. The former body of knowledge is very widely accepted, with every reason to believe it still plays an important role in neurophysiology. One can perhaps suggest that this well-accepted biochemical basis to consciousness may be vestigial, a carry-over from early neural applications in primitive creatures, and that the neurological utilization of the electromagnetic field is a more recent evolutionary strategy, one through which mind first appeared. If this is the case then there may be a bimodal quality to brain function, both biochemical and field-like, with greater use of the latter by higher-level animals, particularly humans.

~~Finally, we cannot avoid commenting that our picture of electromagnetic consciousness viewed as a disembodied entity independent of its constituent elements, also provides a clear alternative to spirit and soul.~~

The author acknowledges K A Jenrow for his helpful discussions and valuable suggestions.

REFERENCES

Alberto D, Busso L, Crotti G, Gandini M, Garfagnini R, Giudici P, Gnesi I, Manta F, and Piragino G, 2008. Effects of static and low-frequency alternating magnetic fields on the ionic electrolytic currents of glutamic acid aqueous solutions. *Electromagn Biol Med.* 27:25-39.

Aspect A, Dalibard J, and Roger G, 1982. Experimental test of Bell's inequalities using time-varying analyzers, *Phys Rev Let* 49: 1804-1807.

Baars B J, and Edelman D B, 2012. Consciousness, biology and quantum hypotheses. *Phys Life Rev* 9: 285-294.

Baureus-Koch C L M, Sommarin M, Persson B R R, Salford L G, and Eberhardt J L, 2003. Interaction between weak low frequency magnetic fields and cell membranes. *Bioelectromagnetics* 24: 395-402.

Bawin S M, and Adey W R, 1976. Sensitivity of calcium binding in cerebral tissue to weak environmental electric fields oscillating at low frequencies. *Proc Natl Acad Sci USA* 73: 1999-2003.

Beason R C, 2005. Mechanisms of magnetic orientation in birds. *Integrat Comp Biol* 45: 565-573.

Blackman C F, Kinney L S, House D E, and Joines W T, 1989. Multiple power-density windows and their possible origin. *Bioelectromagnetics* 10: 115-128.

Blackman C F, Blanchard J P, Benane S G, and House D E, 1994, Empirical test of an ion parametric resonance model for magnetic field interactions with PC-12 cells. *Bioelectromagnetics* 15; 239-250.

Cavagna A, Cimarelli A, Giardina I, Parisi G, Santagati R, Stefanini F, and Viali M, 2010. Scale-free correlations in starling flocks. *Proc Nat Acad Sci USA* 107: 11865-11870.

Chalmers D J, 1995. Facing up to the problem of consciousness. *J Consciousness Studies* 2: 200-219.

Comisso N, Del Giudice E, De Ninno A, Fleischmann M, Giuliani L, Mengoli G, Merlo F, and Talpo G, 2006. Dynamics of the ion resonance effect on amino acids absorbed at the interfaces. *Bioelectromagnetics* 27: 16-25.

Couzin I, 2007. Collective minds. *Nature* 445: 715.

D'Emilio, E, Giuliani L, Lisi A, Ledda M, Grimaldi S, Montagnier L, and Liboff A R, 2014. Lorentz force in water: Evidence that hydronium cyclotron resonance enhances polymorphism. *Electromagn Biol Med* doi:10.3109/15368378.2014.937873.

Fitzsimmons R J, Ryaby J T, Magee F P, and Baylink D J, 1995. Combined magnetic fields increase insulin-like growth factor-II in TE-85 human osteosarcoma bone cell cultures. *Endocrinology* 136: 3100-3106.

Flohr H 2000. NMDA receptor-mediated computational processes and phenomenal consciousness, In: *Neuronal Correlates of Consciousness. Empirical and Conceptual Questions*, Ed. T Metzinger, Cambridge MIT Univ Press, pp 245-258.

Foletti A, Grimaldi S, Lisi A, Ledda M, and Liboff A R, 2013. Bioelectromagnetic medicine: the role of resonance signaling. *Electromag Biology and Med* 32: 484-499.

~~Fourie P C, and Greenwald A A, 2002. The particle swarm optimization algorithm in size and shape optimization. *Struct Multidisc Optim* 23: 259-269.~~

Frilot II C, Carrubba S, and Marino AA, 2014. Sensory transduction of weak electromagnetic fields: role of glutamate neurotransmission mediated by NMDA receptors. *Neurosci* 258: 184-191.

Gaetani R, Ledda M, Barile L, Cimenti I, De Carlo F, Forte E, Ionta V, Giuliani L, D'Emilia E, Frati G, Messina E, Grimaldi S, Giacomella A, and Lisi A, 2009. Differentiation of human adult cardiac stem cells exposed to extremely low-frequency electromagnetic field. *Cardiovasc Res* 82: 411-420.

¹ Goodman R, and Henderson A S, 1988. Exposure of salivary gland cells to low-frequency electromagnetic fields alters polypeptide synthesis. *Proc Natl Acad Sci USA* 85: 3928-3932.

Hameroff S R, 1994. Quantum coherence in microtubules: A neural basis for emergent consciousness? *J Consciousness Studies* 1: 91-118.

Hameroff S, 2012. How quantum brain biology can rescue conscious free will. *Front Integr Neurosci* doi:10.3389/fnint.2012.00093.

¹ Hameroff S, and Penrose R, 2014. Consciousness in the universe: A review of the Orch OR theory. *Phys Life Rev* 11: 39-78.

Hemelrijk C K, and Hildenbrandt H, 2011. Some causes of the variable shape of flocks of birds. *PloS ONE* 6(8):e22479.doi:10.1371/journal.pone.0022479.

Horton P, Ryaby J T, Magee F P, and Weinstein A M, 1993. Stimulation of specific neuronal differentiation proteins in PC-12 cells by combined AC/DC magnetic fields. In Blank, M., Ed., *Electricity and Magnetism in Biology and Medicine*. San Francisco Press, San Francisco.

Jefferys J G R, Traub R D, and Whittington M A, 1996. Neuronal networks for induced “40 Hz” rhythms. *Trends in Neurosciences* 19: 202-208.

Jenrow K A, Zhang X, Renehan W E, and Liboff A R, 1998. Weak ELF magnetic field effects on hippocampal rhythmic slow wave activity. *Exp Neurol* 153: 328-334.

Lednev V V, 1991. Possible mechanism for the influence of weak magnetic fields on biological system. *Bioelectromagnetics* 12: 71-75.

Liboff A R, Williams T Jr, Strong D M, and Wistar R Jr, 1984. Time-varying magnetic fields: effect on DNA synthesis. *Science* 223: 818-820.

Liboff A R, 1985. Geomagnetic cyclotron resonance in living cells. *J Biol Physics* 13: 99-102.

Liboff A R, 1994. The electromagnetic field as a biological variable. In Frey A H, ed, *On the Nature of Electromagnetic Field Interactions with Biological Systems*, R G Landes, Austin.

Liboff A R, 2004. Toward an electromagnetic paradigm for biology and medicine. *J Alt Com Med* 10: 41-47.

Liboff A R, 2005. A Rational Biology, Electromagnetic Biology and Medicine 24: 211-220.

Liboff A R, 2006. The ion cyclotron resonance hypothesis. In: Greenebaum B, Barnes F, editors, Handbook of Bioelectromagnetism, 3rd ed Boca Raton: CRC Press.

Lovely R H, Creim J A, Miller D L, and Anderson L E, 1993. Behavior of rats in a radial arm maze during exposure to magnetic fields: Evidence for effects of magnesium ion resonance. 15th annual meeting *Bioelectromagnetics Society*, Abstract EI-6, Los Angeles.

Lyskov Y B, Chernysev M V, Michailov V O, Kozlov A P, Makarova T M, Vasilyeva Y V, Druzin M Y, Sokolov G V, and Vishnevski A M, 1996. The effect of a magnetic field with the frequency of 50 Hz on behavior in rats depends on the value of the constant magnetic field. *Biophysics* 41: 881-886.

Manikonda P K, Rajendra P, Devandraneth D, Gunasekaran B, Channekeshava, Aradhya R S, Sashidhar R B, and Subramanyam C, 2007. Influence of extremely low frequency magnetic fields on Ca²⁺ signaling and NMDA receptor functions in rat hippocampus. *Neurosci Lett* 413: 145-149.

Marino A A, Nilsen E, Chesson A L, Jr, and Frilot C 2004. Effect of low-frequency magnetic fields on brain electrical activity in human subjects. *Clin Neurophysiol* 115: 1195-1201.

Martin H, and Lindauer M, 1977. The effect of the earth's magnetic field on gravity orientation in the honey bee (*Apis mellifica*). *J Comp Physiol A* 122:145-187.

McFadden J, 2001. Quantum Evolution, pp 295-305, Norton, New York.

McFadden J, 2002. Synchronous firing and its influence on the brain's electromagnetic field. *J Consciousness Studies* 9: 23-50.

McFadden J, 2007. Conscious electromagnetic (CEMI) field theory. *Neuroquantology* 5: 262-270.

McFadden J, 2013. The CEMI field theory: closing the loop. *J Consciousness Studies* 20: 153-168.

Pazur A, 2004. Characterization of weak magnetic field effects in an aqueous glutamic acid solution by nonlinear dielectric spectroscopy and voltammetry. *Biomag Res and Tech* 2: 8. Doi:10.1186/1477-04x-2-8.

Penrose R, 1994. *Shadows of the Mind*, Oxford Univ Press, Oxford.

Penrose R, 1989. *The Emperor's New Mind*, Oxford Univ Press, Oxford.

Pockett S, 2000. *The Nature of Consciousness: A Hypothesis*. Wireless Club Press, New York.

Pockett S, 2011. Initiation of intentional activities and the electromagnetic field theory of consciousness. *Humana Monte* 15: 159-175.

Pockett S, 2012. The electromagnetic field theory of consciousness: a testable hypothesis about the characteristics of conscious as opposed to non-conscious fields. *J Consciousness Studies* 19: 191-223.

Pockett S 2013. Field theories of consciousness. *Scholarpedia* doi:10.4249/Scholarpedia.4951

Smith S D, Liboff A R, McLeod B R, and Barr E J, 1992. Effects of ion resonance tuned magnetic fields on N-18 neuroblastoma cells. In Allen, M.J., Cleary, A.F., Sowers, A.E., and Shillady, D.D., Eds *Charge and Field Effects in Biosystems-3*, Birkhauser, Boston, pp. 263-271.

Takashima S, Onaral B, and Schwan H P, 1979. Effects of modulated RF energy on the EEG of mammalian brains: Effects of acute and chronic irradiations. *Radiation and Environmental Biophysics* 16: 15-27.

Thomas J R, Schrot J, and Liboff A R, 1986. Low-intensity magnetic fields alter operant behavior in rats. *Bioelectromagnetics* 7:349-357.

Tomoni G, and Edelman G M, 1998. Consciousness and complexity. *Science* 282: 1846-1851.

Vorobyov V V, Galchenko A A, Kukushkin N I, and Akoev I G, 1997. Effects of weak microwave fields amplitude modulated at ELF on EEG of symmetric brain areas in rats. *Bioelectromagnetics* 18: 293-298.

Vorobyov V V, Sosunov E A, Kukushkin N I, and Lednev V V, 1998. Weak combined magnetic field affects basic and morphine-induced rat's EEG. *Brain Res* 781: 182-187.

Vorobyov V V, Janac B, Pesic V, and Prolic Z, 2010. Repeated exposure to low-level extremely low frequency-modulated microwaves affects cortex-hypothalamus interplay in freely moving rats: EEG study. *Int J Radiat Biol* 86: 376-383.

Walker M M, and Bitterman M E, 1989.. Honeybees can be trained to respond to very small changes in geomagnetic field intensity. *J Exp Biol* 145: 489-494.

Walleczek J, and Liburdy R P, 1990, Nonthermal 60 Hz sinusoidal magnetic field exposure enhances $^{45}\text{Ca}^{2+}$ uptake in rat thymocytes: dependence on mitogen activation. *FEBS Lett* 271: 157-160.

Zhadin M N, Novikov V V, Barnes F S, and Pergola N F, 1998. Combined action of static and alternating magnetic fields on ionic current in aqueous glutamic acid solutions. *Bioelectromagnetics* 19: 41-45.

Zhadin M N, Deryugina O N, and Pisachenko T M, 1999. Influence of combined DC and AC magnetic fields on rat behavior. *Bioelectromagnetics* 20: 378-386.